随着病情恶化，青光眼患者会逐渐开始失去周边（侧面）视力，而其他人可能会出现中央视野视力损失。



周边（侧面）视力损失



中央视野视力损失

如不及时治疗，可能会导致视力障碍并最终失明。

什么是高眼压症？

- 高眼压症是指眼内液体排出不畅，导致眼压高于正常水平
- 高眼压症通常没有体征或症状，因此确诊患者应由眼科医生密切监测，因为他们可能有更高的青光眼风险

重要安全信息（续）

- 避免让容器尖端接触眼睛或任何其他表面（包括手指），以防止溶液被细菌污染。使用受污染的溶液可能造成眼睛严重损伤，甚至可能导致视力丧失。

请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。

节省治疗旅程费用

持有 Medicare Part D？

VYZULTA 现已覆盖**近 77%*** 持有 Medicare 处方药承保的患者

*基于 2025 年 3 月的数据。高达 76.7% 的 Medicare Part D 患者可获得保障。

符合“Extra Help”计划资格的患者只需支付约 $12 或更少费用即可获得 VYZULTA

如果您参加了 Medicare 和 Medicaid，即已自动符合“Extra Help”计划资格。

符合条件† 且商业保险承保的患者首次处方和符合条件的续药**可能只需支付 $25 共付额即可获得最多 90 天用量**。

仅限参与活动的独立药店。在非参与药店，符合条件且商业保险承保的患者可能只需支付低至 $35 共付额。

†条款、条件和限制适用。详见 BauschAccessProgram.com 上的资格标准和条款及条件。

重要安全信息（续）

- 研究中观察到的最常见副作用包括眼睛发红、眼睛刺激、眼睛疼痛以及滴药后的疼痛感。

如果处方药造成不良副作用，鼓励您向 FDA 报告。请访问 www.fda.gov/medwatch 或致电 1-800-FDA-1088

适应症

VYZULTA®（拉坦前列腺素布诺滴眼液），0.024%，用于降低开角型青光眼或高眼压症患者的眼内压。

BAUSCH + LOMB

©2025 Bausch + Lomb. VYZ.0045.USA.25

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%

SIMPLIFIED CHINESE
简体中文

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%



掌控您的治疗旅程

非真实患者。



您的视力。
您的 VYZULTA。

降低开角型青光眼或高眼压症引起的眼压

*请参阅完整的处方信息。

重要安全信息

- 虹膜可能出现棕色或棕色加深，这种变化可能是永久性的。还可能发生眼睑皮肤变黑，这通常在停止 VYZULTA 治疗后是可逆的。

请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。

了解眼压

什么是开角型青光眼?

- 开角型青光眼是一种逐渐引起视力丧失并可能导致失明的疾病
- 当单眼或双眼的液体排出不畅时就会发生这种情况
- 这会增加眼压,进而可能损害视神经

青光眼通常被称为*隐形的视力窃贼*,因为它直到损伤严重时才会出现症状。这就是为什么早期检测和定期眼科检查对保护视力至关重要。尽管青光眼无法治愈或逆转,但降低眼压可以帮助保护视力。



非真实患者。

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%

VYZULTA 是什么?

- VYZULTA 是一种治疗滴眼液,可能有助于降低开角型青光眼或高眼压症患者的眼压
- 降低眼压可以帮助减少青光眼导致的视力丧失风险

定期进行眼科检查,并按处方每晚使用一次 VYZULTA 以帮助保护您的视力

按处方每天使用 VYZULTA

VYZULTA 可以帮助降低眼压,这对控制您的青光眼或高眼压症很重要。**按指示使用 VYZULTA 可以支持您的治疗目标。**

VYZULTA 尚无通用等效产品。**如果药店建议使用另一种产品,那就不是 VYZULTA。**如果药店缺货,请要求他们从批发商处订购。

担心眼睛发红?

一些青光眼治疗可能引发较高概率的眼部充血症状,从而导致治疗中断。

在临床试验中,**VYZULTA 导致的眼睛充血发红率很低,仅不到 1% 的患者因副作用而停止使用。**

访问 VYZULTA.com 以了解更多

扫描二维码或访问 VYZULTA.com 获取治疗成功秘诀、资源和优惠信息。

重要安全信息(续)

- 睫毛可能逐渐发生变化,包括长度增加、厚度、颜色、数量和/或生长方向改变。这些变化可能导致双眼的外观差异。这些变化通常在停止 VYZULTA 治疗后是可逆的。

重要安全信息(续)

- 如果您的眼睛发生任何变化、新出现任何眼部状况、视力突然变化、出现任何眼部副作用或接受了眼部手术,请立即告知医生。
- 滴 VYZULTA 前请摘取隐形眼镜。15 分钟后方可重新佩戴。

重要安全信息(续)

- 如果您同时使用其他滴眼液和 VYZULTA,请间隔 5 分钟使用每种滴眼液。

请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。

Source segment	Translation	Back Translation
Contents	目录	Table of Contents
©2025 Bausch + Lomb.	©2025 Bausch + Lomb.	©2025 Bausch + Lomb.
VYZ.0045.USA.25	VYZ.0045.USA.25	VYZ.0045.USA.25
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全信息（续）	Important Safety Information (Continued)
The most common side effects seen in studies were eye redness, eye irritation, eye pain, and pain after putting the drops in your eye.	研究中观察到的最常见副作用包括眼睛发红、眼睛刺激、眼睛疼痛以及滴药后的疼痛感。	The most common side effects observed in the studies included eye redness, eye irritation, eye pain, and pain after the drug was applied.
You are encouraged to report negative side effects of prescription drugs to the FDA.	如果处方药造成不良副作用，鼓励您向 FDA 报告。	If a prescription drug causes an adverse side effect, you are encouraged to report it to the FDA.
Visit www.fda.gov/medwatch or call 1-800-FDA-1088	请访问 www.fda.gov/medwatch 或致电 1-800-FDA-1088	Please visit www.fda.gov/medwatch or call 1-800-FDA-1088
INDICATION	适应症	Indications
VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024%, is used to lower intraocular (eye) pressure in patients with open-angle glaucoma or ocular hypertension.	VYZULTA®（拉坦前列腺素布诺滴眼液），0.024%，用于降低开角型青光眼或高眼压症患者的眼内压。	VYZULTA® (Latanoprost Bruno Eye Drops), 0.024%, is used to reduce intraocular pressure in patients with open-angle glaucoma or ocular hypertension.
† Terms, conditions, and limitations apply.	† 条款、条件和限制适用。	† Terms, conditions and restrictions apply.
Please see eligibility criteria and terms and conditions at BauschAccessProgram.com.	详见 BauschAccessProgram.com 上的资格标准和条款及条件。	See eligibility criteria and terms and conditions at BauschAccessProgram.com.
Eligible† commercially insured patients may pay as little as a $25 copay for up to a 90-day supply for first Rx and eligible refills.	符合条件†且商业保险承保的患者首次处方和符合条件的续药可能只需支付 $25 共付额即可获得最多 90 天用量。	Eligible† patients with commercial insurance coverage may only need to pay $25 copay for up to a 90-day supply of their first prescription and eligible refills.
Exclusively at participating independent pharmacies.	仅限参与活动的独立药店。	Participating independent pharmacies only.
At non-participating pharmacies, eligible commercial patients may pay as little as a $35 copay.	在非参与药店，符合条件且商业保险承保的患者可能只需支付低至 $35 共付额。	At non-participating pharmacies, eligible patients with commercial insurance coverage may only need to pay a copay as low as $35.
Those who qualify for the "Extra Help" Program pay about $12 or less for VYZULTA	符合"Extra Help"计划资格的患者只需支付约 $12 或更少费用即可获得 VYZULTA	Patients who qualify for the "Extra Help" program can get VYZULTA for approximately $12 or less
If you are enrolled in Medicare and Medicaid, you already qualify for the "Extra Help" Program	如果您参加了 Medicare 和 Medicaid，即已自动符合"Extra Help"计划资格。	If you have Medicare and Medicaid, you automatically qualify for the "Extra Help" program.
* Based on data from March 2025.	* 基于 2025 年 3 月的数据。	* Based on data as of March 2025.
Up to 76.7% of Medicare Part D patients may be covered.	高达 76.7% 的 Medicare Part D 患者可获得保障。	As high as 76.7% of Medicare Part D patients are covered.
VYZULTA is now covered for nearly 77%* of	VYZULTA 现已覆盖近 77%* 持有 Medicare	VYZULTA is now available for nearly

patients with Medicare prescription coverage	处方药承保的患者	77%*patients with Medicare prescription drug coverage
Have Medicare Part D?	持有 Medicare Part D？	Have Medicare Part D?
TREATMENT JOURNEY SAVINGS	节省治疗旅程费用	Save money on treatment trips
Take control of your treatment journey	掌控您的治疗旅程	Take control of your treatment journey
IMPORTANT SAFETY INFORMATION	重要安全信息	Important Safety Information
Brown color, or increased brown color, of the iris can occur, which may be permanent.	虹膜可能出现棕色或棕色加深，这种变化可能是永久性的。	The iris may appear brown or darken in color, and this change may be permanent.
Darkening of the eyelid skin may also occur, which is usually reversible after VYZULTA treatment is discontinued.	还可能发生眼睑皮肤变黑，这通常在停止 VYZULTA 治疗后是可逆的。	Darkening of the eyelid skin may also occur and is generally reversible after cessation of VYZULTA treatment.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。	Please see additional “Important Safety Information” in the text and full “Prescribing Information” in the pocket.
* See full Prescribing Information.	* 请参阅完整的处方信息。	* Please see full Prescribing Information.
YOUR VISION.	您的视力。	Your vision.
YOUR VYZULTA.	您的 VYZULTA。	Your VYZULTA.
Lower your eye pressure from open-angle glaucoma or ocular hypertension	降低开角型青光眼或高眼压症引起的眼压	Lower intraocular pressure caused by open-angle glaucoma or ocular hypertension
One Drop	每日 一次	Once a day
Once A Day*	每次一滴*	One drop at a time*
Not an actual patient.	非真实患者。	Not a real patient.
SIMPLIFIED CHINESE	SIMPLIFIED CHINESE 简体中文	SIMPLIFIED CHINESE SIMPLIFIED CHINESE
IMPORTANT SAFETY INFORMATION (cont’d)	重要安全信息（续）	Important Safety Information (Continued)
Avoid touching the tip of the container to the eye or any other surface, including a finger, in order to avoid contamination of the solution by bacteria.	避免让容器尖端接触眼睛或任何其他表面（包括手指），以防止溶液被细菌污染。	Avoid letting the container tip touch your eyes or any other surface (including fingers) to prevent contamination of the solution with bacteria.
Serious damage to the eye and possible loss of vision may result from using contaminated solutions.	使用受污染的溶液可能造成眼睛严重损伤，甚至可能导致视力丧失。	Use of contaminated solutions may cause serious eye damage, possibly resulting in loss of vision.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。	Please see additional “Important Safety Information” in the text and full “Prescribing Information” in the pocket.
What is ocular hypertension?	什么是高眼压症？	What is ocular hypertension?
Ocular hypertension is when the fluid in your eye is not draining properly which causes the	高眼压症是指眼内液体排出不畅，导致眼压高于正常水平	Ocular hypertension is a condition in which fluid in the eye is not draining properly, causing eye

pressure in your eye to be higher than normal		pressure to be higher than normal.
There are no signs or symptoms for ocular hypertension, so people diagnosed with this condition should be monitored carefully by an eye doctor, as they may be at a higher risk of glaucoma	高眼压症通常没有体征或症状，因此确诊患者应由眼科医生密切监测，因为他们可能有更高的青光眼风险	Ocular hypertension often has no signs or symptoms, so people diagnosed with it should be closely monitored by an eye doctor as they may be at increased risk for glaucoma.
If left untreated, this may lead to a visual disability and, eventually, blindness.	如不及时治疗，可能会导致视力障碍并最终失明。	If not treated promptly, it may lead to visual impairment and eventually blindness.
Central field vision loss	中央视野视力损失	Central visual field loss
Peripheral (side) vision loss	周边（侧面）视力损失	Peripheral (side) vision loss
As the condition worsens, people with glaucoma will slowly begin to lose peripheral (side) vision, while others may experience central field loss.	随着病情恶化，青光眼患者会逐渐开始失去周边（侧面）视力，而其他人可能会出现中央视野视力损失。	As the disease progresses, people with glaucoma gradually begin to lose peripheral (side) vision, while others may experience central visual field loss.
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全信息（续）	Important Safety Information (Continued)
If you are using other eye drops in addition to VYZULTA, wait 5 minutes between using each type of eye drop.	如果您同时使用其他滴眼液和 VYZULTA，请间隔 5 分钟使用每种滴眼液。	If you are using other eye drops and VYZULTA, take 5 minutes between using each eye drop.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	请参阅文中的附加“重要安全信息”以及手册夹层袋内完整“处方信息”。	Please see additional “Important Safety Information” in the text and full “Prescribing Information” in the pocket.
Learn more at VYZULTA.com	访问 VYZULTA.com 以了解更多	Visit VYZULTA.com to learn more
Scan the QR code or visit VYZULTA.com for tips for treatment success, resources, and savings offers.	扫描二维码或访问 VYZULTA.com 获取治疗成功秘诀、资源和优惠信息。	Scan the QR code or visit VYZULTA.com for treatment success tips, resources and offers.
Concerned about eye redness?	担心眼睛发红？	Worried about red eyes?
Some glaucoma treatments may cause high rates of eye redness, leading to discontinuation.	一些青光眼治疗可能引发较高概率的眼部充血症状，从而导致治疗中断。	Some glaucoma treatments may cause a high incidence of eye redness, which may lead to discontinuation of treatment.
In clinical trials, VYZULTA had low rates of eye redness and <1% of patients stopped using it due to side effects.	在临床试验中，VYZULTA 导致的眼睛充血发红率很低，仅不到 1% 的患者因副作用而停止使用。	In clinical trials, VYZULTA caused very little eye redness, with less than 1% of patients discontinuing use due to side effects.
Use VYZULTA daily as prescribed	按处方每天使用 VYZULTA	Use VYZULTA daily as prescribed
VYZULTA may help lower your eye pressure, which is important in managing your glaucoma or ocular hypertension.	VYZULTA 可以帮助降低眼压，这对控制您的青光眼或高眼压症很重要。	VYZULTA can help lower the pressure in your eye, which is important to control your glaucoma or ocular hypertension.
Using VYZULTA as directed may support your treatment goals.	按指示使用 VYZULTA 可以支持您的治疗目标。	Use VYZULTA as directed to support your treatment goals.

There is no generic equivalent for VYZULTA.	VYZULTA 尚无通用等效产品。	There are no generic equivalents for VYZULTA.
If the pharmacy suggests a different product, it is not the same as VYZULTA.	如果药店建议使用另一种产品，那就不是 VYZULTA。	If the pharmacy recommends another product, it is not VYZULTA.
If the pharmacy is out of stock, ask them to order it from the wholesaler.	如果药店缺货，请要求他们从批发商处订购。	If the pharmacy is out of stock, ask them to order it from a wholesaler.
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全信息（续）	Important Safety Information (Continued)
Gradual changes to eyelashes may occur, including increased length, thickness, color, number of eyelashes and/or direction of eyelash growth.	睫毛可能逐渐发生变化，包括长度增加、厚度、颜色、数量和/或生长方向改变。	Eyelashes may gradually change, including increasing in length, thickness, color, number, and/or changing direction of growth.
These changes may result in a difference in appearance of each eye.	这些变化可能导致双眼的外观差异。	These changes can cause the appearance of the two eyes to differ.
These changes are usually reversible after VYZULTA treatment is discontinued.	这些变化通常在停止 VYZULTA 治疗后是可逆的。	These changes were generally reversible after cessation of VYZULTA treatment.
Glaucoma is often called the silent thief of sight because it shows no symptoms until the damage is advanced.	青光眼通常被称为隐形的视力窃贼，因为它直到损伤严重时才会出现症状。	Glaucoma is often called the invisible thief of vision because it does not cause symptoms until damage is severe.
That's why early detection and regular eye exams are crucial to protecting your vision.	这就是为什么早期检测和定期眼科检查对保护视力至关重要。	That's why early detection and regular eye exams are crucial to protecting your vision.
While glaucoma can't be cured or reversed, lowering eye pressure can help preserve sight.	尽管青光眼无法治愈或逆转，但降低眼压可以帮助保护视力。	Although glaucoma cannot be cured or reversed, lowering eye pressure can help protect your vision.
What is open-angle glaucoma?	什么是开角型青光眼？	What is open-angle glaucoma?
Open-angle glaucoma is a disease that gradually causes vision loss and may lead to blindness	开角型青光眼是一种逐渐引起视力丧失并可能导致失明的疾病	Open-angle glaucoma is a disease that gradually causes vision loss and may lead to blindness.
It occurs when the fluid in one or both of your eyes does not drain properly	当单眼或双眼的液体排出不畅时就会发生这种情况	This happens when fluid doesn't drain properly from one or both eyes.
This increases the pressure in your eye, which may damage the optic nerve	这会增加眼压，进而可能损害视神经	This increases eye pressure, which can damage the optic nerve
UNDERSTANDING EYE PRESSURE	了解眼压	Understanding intraocular pressure
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全信息（续）	Important Safety Information (Continued)
Tell your doctor immediately about any change in your eye, any eye condition that develops, any sudden change in your vision, any eye side effects, or if you have eye surgery.	如果您的眼睛发生任何变化、新出现任何眼部状况、视力突然变化、出现任何眼部副作用或接受了眼部手术，请立即告知医生。	Tell your doctor right away if you have any changes in your eyes, develop any new eye condition, have a sudden change in vision, experience any eye side effects, or have had eye surgery.
Remove contact lenses before putting VYZULTA in your eyes.	滴 VYZULTA 前请摘取隐形眼镜。	Remove contact lenses before using VYZULTA.

Lenses may be reinserted 15 minutes later.	15 分钟后方可重新佩戴。	Wait 15 minutes before putting them back on.
Keep up with regular eye appointments and use VYZULTA once daily in the evening as prescribed to help preserve your vision	定期进行眼科检查，并按处方每晚使用一次 VYZULTA 以帮助保护您的视力	Get regular eye exams and use VYZULTA nightly as prescribed to help protect your vision
What is VYZULTA?	VYZULTA 是什么？	What is VYZULTA?
VYZULTA is a treatment drop that may help reduce eye pressure in people with open-angle glaucoma or ocular hypertension	VYZULTA 是一种治疗滴眼液，可能有助于降低开角型青光眼或高眼压症患者的眼压	VYZULTA is a therapeutic eye drop that may help reduce intraocular pressure in people with open-angle glaucoma or ocular hypertension
Lowering eye pressure can help reduce the risk of vision loss due to glaucoma	降低眼压可以帮助减少青光眼导致的视力丧失风险	Lowering eye pressure can help reduce risk of vision loss from glaucoma
Not an actual patient.	非真实患者。	Not a real patient.
Index	索引	index