隨著病情惡化，青光眼患者會逐漸開始失去週邊（側面）視力，而有些患者則可能會出現中央視野喪失的情況。



週邊（側面）視野喪失



中央視野喪失

若不加以治療，可能會導致視力障礙，最終導致失明。

什麼是眼高壓？

- 眼高壓是指眼內液體無法正常排出，導致眼壓高於正常值的情況
- 眼高壓沒有任何徵兆或症狀，因此被診斷出患有此疾病者應接受眼科醫師的密切監測，因為這類患者罹患青光眼的風險可能更高

重要安全資訊（續）

- 避免讓容器尖端接觸到眼睛或任何其他表面（包括手指），以免溶液受到細菌污染。使用受污染的溶液可能會對眼睛造成嚴重傷害，甚至可能導致視力喪失。

請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。

節省療程費用

您擁有 Medicare Part D 嗎？

VYZULTA 目前涵蓋**近 77%* 享有 Medicare 處方藥保險的患者**

*此為 2025 年三月的資料。高達 76.7% 的 Medicare Part D 患者可獲得保障。

符合「Extra Help」計畫資格者僅需支付約 $12 或更少的費用即可取得 VYZULTA

若您已加入 Medicare 和 Medicaid，便已獲得「Extra Help」計畫使用資格

符合資格†且有商業保險的患者，首次處方和符合資格的續方，**最多可獲得 90 天的藥量供應，自付額可能低至 $25。**

僅限參與活動的獨立藥局。在非參與藥局，符合資格的商業患者可能僅需支付 $35 的自付額。

†適用條款、條件和限制。請參閱 BauschAccessProgram.com 上的資格標準及條款和條件。

重要安全資訊（續）

- 研究發現，最常見的副作用為眼睛發紅、眼睛刺激、眼睛疼痛以及滴眼藥水後感到疼痛。

我們建議您向 FDA 回報處方藥的負面副作用。請造訪 www.fda.gov/medwatch 或致電 1-800-FDA-1088

適用症狀

VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024% 用於降低開角型青光眼或眼高壓患者的眼壓。

BAUSCH + LOMB

©2025 Bausch + Lomb. VYZ.0047.USA.25

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%

TRADITIONAL CHINESE
繁體中文



掌控
您的治療歷程

圖中人物非真實病患。



您的視界。
您的 VYZULTA。

降低因開角型青光眼或眼高壓引起的眼壓

*參閱完整處方資訊。

重要安全資訊

- 虹膜可能會變成棕色，或棕色部分增多，此改變可能是永久性的。眼瞼皮膚也可能變黑，通常在停止使用 VYZULTA 治療後便能恢復。

請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。

了解眼壓

什麼是開角型青光眼？

- 開角型青光眼是一種會逐漸導致視力喪失，並可能導致失明的疾病
- 當一隻或兩隻眼睛裡的液體無法正常排出時，就會引發這種疾病
- 此時眼壓增加，可能會損傷視神經

青光眼通常被稱為*無聲的視力竊賊*，因為在病情發展到較嚴重程度之前，它不會表現出任何症狀。這就是為什麼及早發現和定期眼科檢查對保護視力至關重要。雖然青光眼無法治愈或逆轉，但降低眼壓有助於保護視力。



圖中人物非真實病患。

VYZULTA®
(latanoprostene bunod ophthalmic solution), 0.024%

VYZULTA 是什麼？

- VYZULTA 是一款治療用眼藥水，有助於降低開角型青光眼或眼高壓患者的眼壓
- 降低眼壓有助於降低青光眼導致視力喪失的風險

定期進行眼科檢查，並根據處方每晚使用一次 VYZULTA，以協助保護您的視力

請根據處方每日使用 VYZULTA

VYZULTA 有助於降低您的眼壓，對於控制您的青光眼或眼高壓相當重要。**按照指示使用 VYZULTA 有助於達成您的治療目標。**

VYZULTA 沒有同效的仿製藥。**若藥局建議使用不同產品，則該產品與 VYZULTA 並不相同。**若藥局缺貨，可請藥局向批發商訂購。

擔心用了眼睛發紅？

某些青光眼療程可能會造成高機率的眼睛發紅症狀，進而導致治療中斷。

在臨床試驗中，**VYZULTA 的眼睛發紅發生率極低，因副作用而停止使用的患者人數僅不到 1%。**

請造訪 VYZULTA.com 了解更多資訊

請掃描二維碼或造訪 VYZULTA.com 了解成功治療的訣竅、相關資源以及節省費用的優惠方案。

重要安全資訊（續）

- 睫毛可能會逐漸發生變化，包括睫毛變長、變粗、顏色變深、數量增加和 / 或生長方向改變。這些變化可能會導致兩眼的外觀不同。停止 VYZULTA 治療後，這些變化通常能夠恢復原狀。

重要安全資訊（續）

- 如果您的眼睛出現任何變化、患上任何眼部疾病、視力突然變化、出現任何眼部副作用或者您接受了眼部手術，請立即告知醫師。
- 在向眼內滴入 VYZULTA 之前，請取下隱形眼鏡。15 分鐘後即可戴回。

重要安全資訊（續）

- 若您除了 VYZULTA 之外還有使用其他眼藥水，請在使用每種眼藥水之間等待 5 分鐘。

請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。

Source segment	Translation	Back Translation
Contents	內容	content
©2025 Bausch + Lomb.	©2025 Bausch + Lomb.	©2025 Bausch + Lomb.
VYZ.0047.USA.25	VYZ.0047.USA.25	VYZ.0047.USA.25
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全資訊（續）	Important Safety Information (continued)
The most common side effects seen in studies were eye redness, eye irritation, eye pain, and pain after putting the drops in your eye.	研究發現，最常見的副作用為眼睛發紅、眼睛刺激、眼睛疼痛以及滴眼藥水後感到疼痛。	The study found that the most common side effects were eye redness, eye irritation, eye pain and pain after putting the eye drops in.
You are encouraged to report negative side effects of prescription drugs to the FDA.	我們建議您向 FDA 回報處方藥的負面副作用。	We recommend that you report adverse side effects of the prescription drug to the FDA.
Visit www.fda.gov/medwatch or call 1-800-FDA-1088	請造訪 www.fda.gov/medwatch 或致電 1-800-FDA-1088	Please visit www.fda.gov/medwatch or call 1-800-FDA-1088
INDICATION	適用症狀	Indications
VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024%, is used to lower intraocular (eye) pressure in patients with open-angle glaucoma or ocular hypertension.	VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024% 用於降低開角型青光眼或眼高壓患者的眼壓。	VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024% is used to reduce intraocular pressure in patients with open-angle glaucoma or ocular hypertension.
† Terms, conditions, and limitations apply.	† 適用條款、條件和限制。	† Terms, conditions and restrictions apply.
Please see eligibility criteria and terms and conditions at BauschAccessProgram.com.	請參閱 BauschAccessProgram.com 上的資格標準及條款和條件。	Please see eligibility criteria and terms and conditions at BauschAccessProgram.com.
Eligible† commercially insured patients may pay as little as a $25 copay for up to a 90-day supply for first Rx and eligible refills.	符合資格†且有商業保險的患者，首次處方和符合資格的續方，最多可獲得 90 天的藥量供應，自付額可能低至 $25。	Eligible† patients with commercial insurance may receive up to a 90-day supply for first-time prescriptions and eligible refills with a copay as low as $25.
Exclusively at participating independent pharmacies.	僅限參與活動的獨立藥局。	Participating independent pharmacies only.
At non-participating pharmacies, eligible commercial patients may pay as little as a $35 copay.	在非參與藥局，符合資格的商業患者可能僅需支付 $35 的自付額。	At non-participating pharmacies, eligible commercial patients may only need to pay a $35 copay.
Those who qualify for the "Extra Help" Program pay about $12 or less for VYZULTA	符合「Extra Help」計畫資格者僅需支付約 $12 或更少的費用即可取得 VYZULTA	Those who qualify for the "Extra Help" program can get VYZULTA for about $12 or less.
If you are enrolled in Medicare and Medicaid, you already qualify for the "Extra Help" Program	若您已加入 Medicare 和 Medicaid，便已獲得「Extra Help」計畫使用資格	If you are enrolled in Medicare and Medicaid, you are already eligible for the "Extra Help" program.
* Based on data from March 2025.	* 此為 2025 年三月的資料。	* This data is as of March 2025.
Up to 76.7% of Medicare Part D patients may be covered.	高達 76.7% 的 Medicare Part D 患者可獲得保障。	As high as 76.7% of Medicare Part D patients are covered.
VYZULTA is now covered for nearly 77%* of patients with Medicare prescription coverage	VYZULTA 目前涵蓋近 77%* 享有 Medicare 處方藥保險的患者	VYZULTA currently coversnearly 77%* of patients who have Medicare prescription drug coverage

Have Medicare Part D?	您擁有 Medicare Part D 嗎?	Do you have Medicare Part D?
TREATMENT JOURNEY SAVINGS	節省療程費用	Save treatment costs
Take control of your treatment journey	掌控您的治療歷程	Take control of your treatment journey
IMPORTANT SAFETY INFORMATION	重要安全資訊	Important Safety Information
Brown color, or increased brown color, of the iris can occur, which may be permanent.	虹膜可能會變成棕色，或棕色部分增多，此改變可能是永久性的。	The iris may turn brown or have more brown areas, and this change may be permanent.
Darkening of the eyelid skin may also occur, which is usually reversible after VYZULTA treatment is discontinued.	眼瞼皮膚也可能變黑，通常在停止使用 VYZULTA 治療後便能恢復。	The skin of the eyelids may also darken, which usually restore after treatment with VYZULTA is stopped.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。	Please see additional Important Safety Information throughout the document and see full Prescribing Information in the pocket.
* See full Prescribing Information.	* 參閱完整處方資訊。	* See full Prescribing Information.
YOUR VISION.	您的視界。	Your vision.
YOUR VYZULTA.	您的 VYZULTA。	Your VYZULTA.
Lower your eye pressure from open-angle glaucoma or ocular hypertension	降低因開角型青光眼或眼高壓引起的眼壓	Lowering intraocular pressure due to open-angle glaucoma or ocular hypertension
One Drop	每次一滴	One Drop at a time
Once A Day*	每日一次*	Once a day*
Not an actual patient.	圖中人物非真實病患。	The people in the picture are not real patients.
TRADITIONAL CHINESE	TRADITIONAL CHINESE 繁體中文	TRADITIONAL CHINESE TRADITIONAL CHINESE
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全資訊（續）	Important Safety Information (continued)
Avoid touching the tip of the container to the eye or any other surface, including a finger, in order to avoid contamination of the solution by bacteria.	避免讓容器尖端接觸到眼睛或任何其他表面（包括手指），以免溶液受到細菌污染。	Avoid letting the container tip touch your eyes or any other surface (including fingers) to prevent contamination of the solution with bacteria.
Serious damage to the eye and possible loss of vision may result from using contaminated solutions.	使用受污染的溶液可能會對眼睛造成嚴重傷害，甚至可能導致視力喪失。	Use of contaminated solutions may cause serious damage to the eyes, possibly resulting in loss of vision.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。	Please see additional Important Safety Information throughout the document and see full Prescribing Information in the pocket.
What is ocular hypertension?	什麼是眼高壓？	What is ocular hypertension?
Ocular hypertension is when the fluid in your eye is not draining properly which causes the pressure in your eye to be higher than normal	眼高壓是指眼內液體無法正常排出，導致眼壓高於正常值的情況	Ocular hypertension is a condition in which fluid in the eye cannot drain properly, causing the eye pressure to be higher than normal.
There are no signs or symptoms for ocular	眼高壓沒有任何徵兆或症狀，因此被診斷出患有	Ocular hypertension has no signs or symptoms,

hypertension, so people diagnosed with this condition should be monitored carefully by an eye doctor, as they may be at a higher risk of glaucoma	此疾病者應接受眼科醫師的密切監測，因為這類患者罹患青光眼的風險可能更高	so people diagnosed with this condition should be closely monitored by an eye doctor because they may be at a higher risk of developing glaucoma.
If left untreated, this may lead to a visual disability and, eventually, blindness.	若不加以治療，可能會導致視力障礙，最終導致失明。	If left untreated, it can lead to visual impairment and eventually blindness.
Central field vision loss	中央視野喪失	Central visual field loss
Peripheral (side) vision loss	週邊（側面）視野喪失	Peripheral (side) vision loss
As the condition worsens, people with glaucoma will slowly begin to lose peripheral (side) vision, while others may experience central field loss.	隨著病情惡化，青光眼患者會逐漸開始失去週邊（側面）視力，而有些患者則可能會出現中央視野喪失的情況。	As the disease progresses, people with glaucoma gradually begin to lose peripheral (side) vision, and some people may experience central visual field loss
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全資訊（續）	Important Safety Information (continued)
If you are using other eye drops in addition to VYZULTA, wait 5 minutes between using each type of eye drop.	若您除了 VYZULTA 之外還有使用其他眼藥水，請在使用每種眼藥水之間等待 5 分鐘。	If you are using other eye drops in addition to VYZULTA, wait 5 minutes between using each eye drop.
Please see additional Important Safety Information throughout and full Prescribing Information in pocket.	請參閱全文中的其他重要安全資訊，並參閱摺頁內之完整處方資訊。	Please see additional Important Safety Information throughout the document and see full Prescribing Information in the pocket.
Learn more at VYZULTA.com	請造訪 VYZULTA.com 了解更多資訊	For more information, please visit VYZULTA.com
Scan the QR code or visit VYZULTA.com for tips for treatment success, resources, and savings offers.	請掃描二維碼或造訪 VYZULTA.com 了解成功治療的訣竅、相關資源以及節省費用的優惠方案。	Scan the QR code or visit VYZULTA.com for tips for successful treatment, resources, and cost-saving offers.
Concerned about eye redness?	擔心用了眼睛發紅？	Worried about red eyes?
Some glaucoma treatments may cause high rates of eye redness, leading to discontinuation.	某些青光眼療程可能會造成高機率的眼睛發紅症狀，進而導致治療中斷。	Some glaucoma treatments may result in a high rate of eye redness, which may lead to treatment interruption.
In clinical trials, VYZULTA had low rates of eye redness and <1% of patients stopped using it due to side effects.	在臨床試驗中，VYZULTA 的眼睛發紅發生率極低，因副作用而停止使用的患者人數僅不到 1%。	In clinical trials,VYZULTA had a very low incidence of eye redness, with less than 1% of patients discontinuing use due to side effects.
Use VYZULTA daily as prescribed	請根據處方每日使用 VYZULTA	Use VYZULTA daily as prescribed
VYZULTA may help lower your eye pressure, which is important in managing your glaucoma or ocular hypertension.	VYZULTA 有助於降低您的眼壓，對於控制您的青光眼或眼高壓相當重要。	VYZULTA helps lower the pressure in your eye, which is important for managing your glaucoma or high pressure in the eye.
Using VYZULTA as directed may support your treatment goals.	按照指示使用 VYZULTA 有助於達成您的治療目標。	Using VYZULTA as directed can help you achieve your treatment goals.
There is no generic equivalent for VYZULTA.	VYZULTA 沒有同效的仿製藥。	There are no generic equivalents of VYZULTA.
If the pharmacy suggests a different product, it is	若藥局建議使用不同產品，則該產品與 VYZULTA	If your pharmacy recommends a different

not the same as VYZULTA.	並不相同。	product, that product is not the same as VYZULTA.
If the pharmacy is out of stock, ask them to order it from the wholesaler.	若藥局缺貨，可請藥局向批發商訂購。	If the pharmacy is out of stock, you can ask the pharmacy to order it from the wholesaler.
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全資訊（續）	Important Safety Information (continued)
Gradual changes to eyelashes may occur, including increased length, thickness, color, number of eyelashes and/or direction of eyelash growth.	睫毛可能會逐漸發生變化，包括睫毛變長、變粗、顏色變深、數量增加和 / 或生長方向改變。	Eyelashes may gradually change, including becoming longer, thicker, darker, more numerous, and/or changing the direction of their growth.
These changes may result in a difference in appearance of each eye.	這些變化可能會導致兩眼的外觀不同。	These changes may cause the two eyes to look different.
These changes are usually reversible after VYZULTA treatment is discontinued.	停止 VYZULTA 治療後，這些變化通常能夠恢復原狀。	These changes usually restore after you stop using VYZULTA.
Glaucoma is often called the silent thief of sight because it shows no symptoms until the damage is advanced.	青光眼通常被稱為無聲的視力竊賊，因為在病情發展到較嚴重程度之前，它不會表現出任何症狀。	Glaucoma is often called the silent thief of sight because it does not cause any symptoms until the disease has progressed to a more severe stage.
That's why early detection and regular eye exams are crucial to protecting your vision.	這就是為什麼及早發現和定期眼科檢查對保護視力至關重要。	That's why early detection and regular eye exams are crucial to protecting your vision.
While glaucoma can't be cured or reversed, lowering eye pressure can help preserve sight.	雖然青光眼無法治愈或逆轉，但降低眼壓有助於保護視力。	While there is no cure or reversal from glaucoma, lowering eye pressure can help protect your vision.
What is open-angle glaucoma?	什麼是開角型青光眼？	What is open-angle glaucoma?
Open-angle glaucoma is a disease that gradually causes vision loss and may lead to blindness	開角型青光眼是一種會逐漸導致視力喪失，並可能導致失明的疾病	Open-angle glaucoma is a disease that gradually causes vision loss and may lead to blindness.
It occurs when the fluid in one or both of your eyes does not drain properly	當一隻或兩隻眼睛裡的液體無法正常排出時，就會引發這種疾病	This condition occurs when fluid in one or both eyes does not drain properly.
This increases the pressure in your eye, which may damage the optic nerve	此時眼壓增加，可能會損傷視神經	Increased intraocular pressure may damage the optic nerve
UNDERSTANDING EYE PRESSURE	了解眼壓	Understanding intraocular pressure
IMPORTANT SAFETY INFORMATION (cont'd)	重要安全資訊（續）	Important Safety Information (continued)
Tell your doctor immediately about any change in your eye, any eye condition that develops, any sudden change in your vision, any eye side effects, or if you have eye surgery.	如果您的眼睛出現任何變化、患上任何眼部疾病、視力突然變化、出現任何眼部副作用或者您接受了眼部手術，請立即告知醫師。	Tell your doctor right away if you have any changes in your eyes, develop any eye disease, have a sudden change in vision, experience any eye side effects, or if you have eye surgery.
Remove contact lenses before putting VYZULTA in your eyes.	在向眼內滴入 VYZULTA 之前，請取下隱形眼鏡。	Remove contact lenses before putting VYZULTA in your eyes.
Lenses may be reinserted 15 minutes later.	15 分鐘後即可戴回。	You can put them back on after 15 minutes.

Keep up with regular eye appointments and use VYZULTA once daily in the evening as prescribed to help preserve your vision	定期進行眼科檢查，並根據處方每晚使用一次VYZULTA，以協助保護您的視力	Get regular eye exams and use VYZULTA nightly as prescribed to help protect your vision
What is VYZULTA?	VYZULTA 是什麼？	What is VYZULTA?
VYZULTA is a treatment drop that may help reduce eye pressure in people with open-angle glaucoma or ocular hypertension	VYZULTA 是一款治療用眼藥水，有助於降低開角型青光眼或眼高壓患者的眼壓	VYZULTA is a therapeutic eye drop that helps reduce intraocular pressure in patients with open-angle glaucoma or ocular hypertension
Lowering eye pressure can help reduce the risk of vision loss due to glaucoma	降低眼壓有助於降低青光眼導致視力喪失的風險	Lowering eye pressure can help reduce the risk of vision loss from glaucoma
Not an actual patient.	圖中人物非真實病患。	The people in the picture are not real patients.
Index	索引	index